EXCEL

A Prospective, Randomized Trial Comparing Everolimus-Eluting Stents and Bypass Graft Surgery in Selected Patients with Left Main Coronary Artery Disease

Gregg W. Stone MD

Joseph F. Sabik, Patrick W. Serruys, Charles A. Simonton, Philippe Généreux, John Puskas, David E. Kandzari, Marie-Claude Morice, Nicholas Lembo, W. Morris Brown, III, David P. Taggart, Adrian Banning, Béla Merkely, Ferenc Horkay, Piet W. Boonstra, Ad Johannes van Boven, Imre Ungi, Gabor Bogáts, Samer Mansour, Nicolas Noiseux, Manel Sabaté, Jose Pomar, Mark Hickey, Anthony Gershlick, Pawel Buszman, Andrzej Bochenek, Erick Schampaert, Pierre Pagé, Ovidiu Dressler, Ioanna Kosmidou, Roxana Mehran, Stuart J. Pocock, and Arie Pieter Kappetein, for the EXCEL Trial Investigators

Disclosures

Gregg W. Stone

None

Study Design



Follow-up: 1 month, 6 months, 1 year, annually through 5 years

Primary endpoint: Measured at a median 3-yr FU, minimum 2-yr FU

Primary Endpoint

Death, Stroke or MI at 3 Years



No. at Risk:

PCI	948	896	875	850	784	445
CABG	957	868	836	817	763	458

Adjudicated Outcomes at 30 Days

	PCI (n=948)	CABG (n=957)	HR [95%CI]	P-value
Death, stroke or MI	4.9%	7.9%	0.61 [0.42, 0.88]	0.008
- Death	1.0%	1.1%	0.90 [0.37, 2.22]	0.82
- Stroke	0.6%	1.3%	0.50 [0.19, 1.33]	0.15
- MI	3.9%	6.2%	0.63 [0.42, 0.95]	0.02
- Peri-procedural	3.6%	5.9%	0.61 [0.40, 0.93]	0.02
- Spontaneous	0.3%	0.3%	1.00 [0.20, 4.95]	1.00
- STEMI	0.7%	2.3%	0.32 [0.14, 0.74]	0.005
- Non-STEMI	3.2%	3.9%	0.82 [0.50, 1.32]	0.41
Death, stroke, MI or IDR	4.9%	8.4%	0.57 [0.40, 0.82]	0.002
- Ischemia-driven revasc (IDR)	0.6%	1.4%	0.46 [0.18, 1.21]	0.11
Stent thrombosis, def/prob	0.6%	0.0%	-	0.01
Graft occlusion, symptomatic	0.0%	1.2%	-	<0.001
Definite stent thrombosis or symptomatic graft occlusion	0.3%	1.2%	0.27 [0.08, 0.97]	0.03

Major Adverse Events Within 30 Days

	PCI (n=948)	CABG (n=957)	RR [95%CI]	P-value
Peri-procedural MAE, any	8.1%	23.0%	0.35 [0.28, 0.45]	<0.001
- Death*	0.9%	1.0%	0.91 [0.39, 2.23]	0.83
- Stroke*	0.6%	1.3%	0.50 [0.19, 1.34]	0.16
- Myocardial infarction*	3.9%	6.2%	0.63 [0.42, 0.95]	0.02
- Ischemia-driven revascularization*	0.6%	1.4%	0.47 [0.18, 1.22]	0.11
- TIMI major/minor bleeding	3.7%	8.9%	0.42 [0.28, 0.61]	<0.001
- Transfusion ≥2 units	4.0%	17.0%	0.24 [0.17, 0.33]	<0.001
- Major arrhythmia**	2.1%	16.1%	0.13 [0.08, 0.21]	<0.001
- Surgery/radiologic procedure	1.3%	4.1%	0.31 [0.16, 0.59]	<0.001
- Renal failure†	0.6%	2.5%	0.25 [0.10, 0.61]	<0.001
- Sternal wound dehiscence	0.0%	2.0%	0.03 [0.00, 0.43]	<0.001
- Infection requiring antibiotics	2.5%	13.6%	0.18 [0.12, 0.28]	<0.001
- Prolonged intubation (>48 hours)	0.4%	2.9%	0.14 [0.05, 0.41]	<0.001
- Post-pericardiotomy syndrome	0.0%	0.4%	0.11 [0.01, 2.08]	0.12

*Adjudicated events; others are site-reported. **SVT requiring cardioversion, VT or VF requiring treatment, or bradyarrhythmia requiring temporary or permanent pacemaker.
†Serum creatinine increased by ≥0.5 mg/dL from baseline or need for dialysis.

Adjudicated Outcomes at 3 Years (i)

	PCI (n=948)	CABG (n=957)	HR [95%CI]	P-value
Death, stroke or MI (1˚ endpoint)	15.4%	14.7%	1.00 [0.79, 1.26]	0.98
- Death	8.2%	5.9%	1.34 [0.94, 1.91]	0.11
- Definite cardiovascular	3.7%	3.4%	1.10 [0.67, 1.80]	0.71
- Definite non-cardiovascular	3.9%	2.3%	1.60 [0.91, 2.80]	0.10
- Undetermined cause	0.8%	0.3%	2.00 [0.50, 7.98]	0.32
- Stroke	2.3%	2.9%	0.77 [0.43, 1.37]	0.37
- MI	8.0%	8.3%	0.93 [0.67, 1.28]	0.64
- Peri-procedural	3.8%	6.0%	0.63 [0.42, 0.96]	0.03
- Spontaneous	4.3%	2.7%	1.60 [0.95, 2.70]	0.07
- STEMI	1.3%	2.8%	0.46 [0.23, 0.91]	0.02
- Non-STEMI	7.0%	5.9%	1.15 [0.80, 1.65]	0.46

Adjudicated Outcomes at 3 Years (ii)

	PCI (n=948)	CABG (n=957)	HR [95%CI]	P-value
Death, stroke, MI or IDR	23.1%	19.1%	1.18 [0.97, 1.45]	0.10
- Ischemia-driven revasc (IDR)	12.6%	7.5%	1.72 [1.27, 2.33]	<0.001
- PCI	10.3%	6.8%	1.57 [1.13, 2.18]	0.006
- CABG	3.5%	0.8%	4.29 [1.88, 9.77]	<0.001
All revascularization	12.9%	7.6%	1.72 [1.27, 2.33]	<0.001
Stent thrombosis, def/prob	1.3%	0.0%	-	<0.001
- Definite	0.7%	0.0%	-	0.01
- Probable	0.7%	0.0%	-	0.01
- Early (0 - 30 days)	0.7%	0.0%	-	0.008
- Late (30 days – 1 year)	0.1%	0.0%	-	0.32
- Very late (1 year - 3 years)	0.5%	0.0%	-	0.05
Graft occlusion, symptomatic	0.0%	5.4%	-	<0.001
Definite stent thrombosis or symptomatic graft occlusion	0.7%	5.4%	0.12 [0.05, 0.28]	<0.001

Conclusions

- Treatment of patients with LMCAD and low or intermediate SYNTAX scores with CoCr-EES resulted in similar rates of the primary endpoint of death, stroke or MI at 3 years, with fewer adverse events within 30 days compared to CABG
- PCI may thus be considered an acceptable or even preferred revascularization modality for selected patients with LMCAD, a decision which should be made after heart team discussion, taking into account each patient's individual circumstances and preferences